

On continuous symmetries of second-order homogeneous linear ordinary differential equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys.: Conf. Ser. 128 012057

(<http://iopscience.iop.org/1742-6596/128/1/012057>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 152.74.220.241

The article was downloaded on 31/08/2012 at 17:52

Please note that [terms and conditions apply](#).

On continuous symmetries of second-order homogeneous linear ordinary differential equations

Adelio R Matamala¹, Cristian A Salas², Marcelo J Alid^{2,3}

¹Facultad de Ciencias Químicas, Universidad de Concepción, Casilla, 160-C, Concepción, Chile.

²Departamento de Física, Universidad de Concepción, Casilla, 160-C, Concepción, Chile.

³Departamento de Matemáticas, Estadística y Física, Universidad de Las Américas, Chile.

E-mail: amatamal@udec.cl

Abstract. In this work, a method to extract continuous symmetries of general second-order linear ordinary differential equation is presented. The formalism is illustrated by two examples.

1. Introduction

Ordinary differential equations (ODEs) appear in many fields of Physics [1] and the analysis of their symmetries plays an important role to extract information about the solutions of those equations [2-4]. In this work, we have applied the Anderson-Kumei-Wulfman method [5-8] to extract continuous symmetries of general second-order linear ordinary differential equation.

In general, we consider general homogeneous linear ODEs represented by the action of the differential operator $\hat{A}(x)$ on a function $f(x)$

$$\hat{A}(x)f(x) = 0, \quad (1)$$

where the differential operator $\hat{A}(x)$ is give by

$$\hat{A}(x) = \alpha_0(x) + \alpha_1(x)\partial_x + \alpha_2(x)\partial_{xx} + \dots \quad (2)$$

In this work we are interested in second order ODEs, then $\alpha_i = 0$, for $i \geq 3$.

A symmetry operator \hat{Q} of equation (1) is defined as a differential operator that maps solution of equation (1) into solution of the same equation, i.e.,

$$\hat{A}(x)g(x) = 0, \quad (3)$$

where

$$g(x) = \hat{Q}(x)f(x). \quad (4)$$

After assuming a particular form for the operators \hat{Q} , condition (3) under constrain (1) defines the symmetries of the ODE (1).

2. Symmetry extraction

Let us start by considering a general second-order homogeneous linear ODE:

$$\frac{d^2y}{dx^2} + p(x)\frac{dy}{dx} + q(x)y(x) = 0. \quad (5)$$

After substituting

$$y(x) = f(x) \exp \left[-\frac{1}{2} \int^x p(\xi) d\xi \right] \quad (6)$$

in equation (5), it is easy to see that the function $f(x)$ satisfies the following differential equation:

$$\frac{d^2f}{dx^2} + v(x)f(x) = 0, \quad (7)$$

where

$$v(x) = q(x) - \frac{1}{4}p^2(x) - \frac{1}{2}\frac{dp}{dx}. \quad (8)$$

We are interested in continuous symmetry generators of the form

$$\hat{Q}(x) = \alpha(x) + \beta(x)\frac{d}{dx} \quad (9)$$

for the equation (7). In other words, if $f(x)$ is a solution of equation (7) then we need to find the functions $\alpha(x)$ and $\beta(x)$ so that:

$$g(x) = \hat{Q}(x)f(x) \quad (10)$$

is also a solution of equation (7), i.e.

$$\frac{d^2g}{dx^2} + v(x)g(x) = 0. \quad (11)$$

Then differentiating equation (10), replacing in (11) and using that $f(x)$ is lineally independent, we obtain the following equation system:

$$\alpha'' - v'\beta - 2v\beta' = 0, \quad (12)$$

$$2\alpha' + \beta'' = 0. \quad (13)$$

The above system determines the conditions for $\alpha(x)$ and $\beta(x)$ function.

Working on equations (12) and (13), we obtain:

$$\beta''' + 4v\beta' + 2v'\beta = 0. \quad (14)$$

Equation (14) may be solved if $v(x)$ function is known.

Let us assume that we know two particular solutions $u_1(x)$ and $u_2(x)$ for equation (7), i.e.

$$u_1'' + vu_1 = 0 \quad \text{and} \quad u_2'' + vu_2 = 0. \quad (15)$$

Now, if we consider the following definition

$$\phi(x) = C_1u_1^2 + C_2u_2^2 + C_3u_1u_2, \quad (16)$$

It is easy to show that $\phi(x)$ function satisfies the equation:

$$\phi''' + 4v\phi' + 2v'\phi = 0. \quad (17)$$

Therefore, equation (14) defines the $\beta(x)$ function as

$$\beta(x) = C_1 u_1^2 + C_2 u_2^2 + C_3 u_1 u_2, \quad (18)$$

where the functions $u_1(x)$ and $u_2(x)$ are two linearly independent particular solutions of equation (7):

$$u_1'' + v u_1 = 0 \quad \text{and} \quad u_2'' + v u_2 = 0.$$

However, it is only necessary to know one solution since the second linear independent solution of equation (7) is obtained by the relation

$$u_2(x) = u_1(x) \int^x \frac{d\xi}{[u_1(\xi)]^2}. \quad (19)$$

Finally, having $\beta(x)$ it is possible to obtain $\alpha(x)$ expansion coefficient by:

$$\alpha'' = -\frac{1}{2}\beta'''. \quad (20)$$

3. Examples

The symmetry extraction method will be exemplified by two simple equations:

3.1. Example 1

Let us consider the simplest second order ODE:

$$f_{xx} = 0. \quad (21)$$

Step 1: Find a particular solution of (21)

$$u_1 = C_1 x. \quad (22)$$

Step 2: Use equation (19), to build a second independent solution

$$u_2 = C_1 x \int^x \frac{d\xi}{(C_1 \xi)^2} = C_2. \quad (23)$$

Step 3: Use equation (18) to obtain $\beta(x)$ function

$$\beta(x) = B_1 x^2 + B_2 + B_3 x. \quad (24)$$

Step 4: Use equation (13) to obtain $\alpha(x)$ function

$$\alpha(x) = -\frac{1}{2} \int^x \beta''(\xi) d\xi = -B_1 x. \quad (25)$$

Step 5: Build the symmetry generators

$$\hat{Q}_1 = \frac{d}{dx}, \quad \hat{Q}_2 = x \frac{d}{dx}, \quad \hat{Q}_3 = x - x^2 \frac{d}{dx}. \quad (26)$$

Step 6: Check the symmetry property

$$\hat{A} \hat{Q}_1 f = 0, \quad \hat{A} \hat{Q}_2 f = 0, \quad \hat{A} \hat{Q}_3 f = 0 \quad (27)$$

Step 7: Find the algebra

$$[\hat{Q}_1, \hat{Q}_2] = \hat{Q}_1, \quad [\hat{Q}_1, \hat{Q}_3] = I - 2\hat{Q}_2, \quad [\hat{Q}_2, \hat{Q}_3] = \hat{Q}_3. \quad (28)$$

And, introducing the new definition:

$$\hat{A}_0 = \hat{Q}_2 - 1/2, \quad \hat{A}_- = \hat{Q}_1, \quad \hat{A}_+ = \hat{Q}_3, \quad (29)$$

the following commutation relations are obtained:

$$[\hat{A}_0, \hat{A}_\pm] = \pm \hat{A}_\pm, \quad [\hat{A}_+, \hat{A}_-] = -2\hat{A}_0. \quad (30)$$

3.1.1. Symmetry Visualization: With the symmetry generators we can obtain the action of this generators on the solution of the original ODE.

The general solution of ODE (21) is:

$$y = a + bx, \tag{31}$$

where a and b are arbitrary constants. Then the actions of \hat{A}_+ , \hat{A}_- and \hat{A}_0 over (31) are

$$\begin{aligned} \exp(\theta\hat{A}_0)(a + bx) &= \bar{a} + \bar{b}x, & \bar{a} &= e^{-\theta/2}a, & \bar{b} &= e^{\theta/2}b \\ \exp(\theta\hat{A}_-)(a + bx) &= \bar{a} + bx, & \bar{a} &= a + \theta b \\ \exp(\theta\hat{A}_+)(a + bx) &= a + \bar{b}x, & \bar{b} &= b + \theta a \end{aligned}$$

Figure 1 shows the plot of (31) for particular values of a and b constants. Figures 2, 3 and 4 show the action of \hat{A}_+ , \hat{A}_- and \hat{A}_0 symmetry operators on solution (31) respectively.




Figure 1




Figure 2




Figure 3




Figure 4

3.2. Example 2

Let us consider the second order ODE:

$$f_{xx} + k^2 f = 0. \tag{32}$$

Step 1: Find a particular solution of (32)

$$u_1 = \cos(kx). \tag{33}$$

Step 2: Use equation (19) to build a second independent solution

$$u_2 = \cos(kx) \int^x \frac{d\xi}{[\cos(k\xi)]^2} = \frac{\sin(kx)}{k}. \tag{34}$$

Step 3: Use equation (18) to obtain $\beta(x)$ function

$$\beta(x) = C_1 \cos^2(kx) + C_2 \frac{\sin^2(kx)}{k^2} + C_3 \frac{\sin(kx)}{k} \cos(kx). \quad (35)$$

Step 4: Use equation (13) to obtain $\alpha(x)$ function

$$\alpha(x) = C_1 \frac{k \sin(2kx)}{2} - C_2 \frac{\sin(2kx)}{2k} + C_3 \frac{1}{2} (1 - \cos(2kx)) + C_4. \quad (36)$$

Step 5: Build the symmetry generators

$$\hat{Q}_1 = \frac{k}{2} \sin(2kx) + \frac{1}{2} (1 + \cos(2kx)) \frac{d}{dx}, \quad (37)$$

$$\hat{Q}_2 = \frac{1}{2} - \frac{1}{2} \cos(2kx) + \frac{1}{2k} \sin(2kx) \frac{d}{dx}, \quad (38)$$

$$\hat{Q}_3 = \frac{1}{2k} \sin(2kx) + \frac{1}{2k^2} (\cos(2kx) - 1) \frac{d}{dx}. \quad (39)$$

Step 6: Check the symmetry property

$$\hat{A} \hat{Q}_1 f = 0 \quad \hat{A} \hat{Q}_2 f = 0, \quad \hat{A} \hat{Q}_3 f = 0. \quad (40)$$

Step 7: Find the algebra

$$[\hat{Q}_1, \hat{Q}_2] = \hat{Q}_1, \quad [\hat{Q}_1, \hat{Q}_3] = I - 2\hat{Q}_2, \quad [\hat{Q}_2, \hat{Q}_3] = \hat{Q}_3. \quad (41)$$

And, introducing the new definitions:

$$\hat{A}_0 = \hat{Q}_2 - 1/2, \quad \hat{A}_- = \hat{Q}_1, \quad \hat{A}_+ = \hat{Q}_3, \quad (42)$$

the following commutation relations are obtained:

$$[\hat{A}_0, \hat{A}_\pm] = \pm \hat{A}_\pm, \quad [\hat{A}_+, \hat{A}_-] = -2\hat{A}_0. \quad (43)$$

3.2.1. Matching examples 1 and 2: From the equation (33) and (34) we have the general solution of EDO (32)

$$y = C_1 \cos(kx) + \frac{C_2}{k} \sin(kx). \quad (44)$$

Taking the limit $k \rightarrow 0$ on equation (32), we obtain equation (21). In same way, this limit reduces solution (44) to the solution of equation (21).

$$y = C_1 + C_2 x. \quad (45)$$

On the other hand, taking the limit $k \rightarrow 0$ on symmetry generators(37), (38) and (39), we obtain

$$\begin{aligned} \lim_{k \rightarrow 0} (\hat{Q}_1) &= \lim_{k \rightarrow 0} \left(\frac{k}{2} \sin(2kx) + \frac{1}{2} (1 + \cos(2kx)) \frac{d}{dx} \right) = \frac{d}{dx}, \\ \lim_{k \rightarrow 0} (\hat{Q}_2) &= \lim_{k \rightarrow 0} \left(\frac{1}{2} - \frac{1}{2} \cos(2kx) + \frac{1}{2k} \sin(2kx) \frac{d}{dx} \right) = x \frac{d}{dx}, \\ \lim_{k \rightarrow 0} (\hat{Q}_3) &= \lim_{k \rightarrow 0} \left(\frac{1}{2k} \sin(2kx) + \frac{1}{2k^2} (\cos(2kx) - 1) \frac{d}{dx} \right) = x - x^2 \frac{d}{dx}. \end{aligned}$$

Showing the compatibility between the symmetry operators obtained in example 1 and 2 respectively. It is noteworthy that commutations relations (28) remain invariant under the limit $k \rightarrow 0$.

4. Conclusion

A general method to find continuous symmetries of second-order linear ODEs has been presented. To obtain the symmetry generators a particular solution of the ODE under study is required. The method has been illustrated by two examples.

Acknowledgments

The authors thank the organizers of 5th International Symposium on Quantum Theory and Symmetries, (Valladolid, Spain), for their warm hospitality. A.R.M. thanks support from FONDECYT (Chile) under Grant No 1050536. Also C.A.S. thanks support from MECESUP (Chile) program under Grant No UCO-0209.

References

- [1] Morse P M and Feshbach H 1953 *Methods of Theoretical Physics. Part I and II*(New York: McGraw-Hill)
- [2] Ibragimov N H 1999 *Elementary Lie Group Analysis and Ordinary Differential Equations*(New York: John Wiley and Sons)
- [3] Hydon P E 2000 *Symmetry Methods for Differential Equations*(New York: Cambridge University Press)
- [4] Cantwell B J 2002 *Introduction to Symmetry Analysis*(New York: Cambridge University Press)
- [5] Anderson R L, Kumei S, and Wulfman C E 1972 *Rev. Mex. Fis.* **21** 1-33
- [6] Anderson R L, Kumei S, and Wulfman C E 1972 *Rev. Mex. Fis.* **21** 35-57
- [7] Anderson R L, Kumei S, and Wulfman C E 1972 *Phys. Rev. Lett.* **28** 988-991
- [8] Anderson R L, Kumei S, and Wulfman C E 1973 *J. Math. Phys.* **14** 1527-1531